arXiv:hep-th/0401005v2 10 Feb 2004

IFT-UAM/CSIC-03-53
hep-th/0401005
January 2nd, 2004

Gödel Spacetimes, Abelian Instantons, the Graviphoton Background and other Flacuum Solutions

Patrick Meessen[1] **and Tomás Ortín**♠ [2]

Instituto de Física Teórica, C-XVI, Universidad Autónoma de Madrid
E-28049-Madrid, Spain

Abstract

We study the relations between all the vacua of Lorentzian and Euclidean $d = 4, 5, 6$ SUGRAs with 8 supercharges, finding a new limiting procedure that takes us from the over-rotating near-horizon BMPV black hole to the Gödel spacetime. The timelike compactification of the maximally supersymmetric Gödel solution of $N = 1, d = 5$ SUGRA gives a maximally supersymmetric solution of pure Euclidean $N = 2, d = 4$ with flat space but non-trivial anti-selfdual vector field flux (*flacuum*) that, on the one hand, can be interpreted as an $U(1)$ instanton on T^4 and that, on the other hand, coincides with the graviphoton background shown by Berkovits and Seiberg to produce the C-deformation introduced recently by Ooguri and Vafa. We construct flacuum solutions in other theories such as Euclidean type IIA supergravity.

---

[1] E-mail: `Patrick.Meessen@uam.es`
[2] E-mail: `Tomas.Ortin@cern.ch`

Contents

1 Introduction **2**

2 Dimensional reduction of $N = (1, 0), d = 6$ SUGRA in the timelike direction **5**

3 Dimensional reduction of $N = 1, \hat{d} = 5$ SUGRA in the timelike direction **7**

4 Maximally supersymmetric vacua of Euclidean $N = 1, d = 5$ SUGRA **8**
4.1 The $aDS_3 \times S^3$ vacuum . . . 9
4.2 The $KG6$ Hpp-wave vacuum . . . 11

5 Maximally supersymmetric vacua of Euclidean $N = 2_\pm, d = 4$ SUGRA **12**
5.1 NH limits of the extreme critically rotating and over-rotating BMPV BHs . 13
5.2 The Euclidean NH limits of extreme rotating BMPV BHs . . . 13
5.3 The Euclidean and Lorentzian Gödel solutions and the flacuum solution . . 13
5.4 The flacuum solution as an Abelian instanton . . . 16

6 (Anti-) selfdual vacua in Euclidean SUGRA **18**
6.1 pp-waves, lightlike reductions and flacuum solutions . . . 21

7 Limiting Solutions **22**
7.1 The limit of the Euclidean Dyonic Robinson-Bertotti solutions . . . 23
7.2 Boosted Hopf oxidation of $aDS_5 \times S^5$ . . . 23

8 Conclusions **24**

A The detailed dimensional reductions **26**
A.1 $N = (1, 0), \hat{d} = 6$ SUGRA . . . 26
A.2 $N = 1, \hat{d} = 5$ SUGRA . . . 28

1 Introduction

There has been much progress recently in the classification of supersymmetric vacua in $d = 4, 5, 6$ Lorentzian supergravities (SUGRAs) with eight supercharges [1, 2, 3, 4] and in the study of their relations via dimensional reduction and oxidation [5, 9]. The study of these theories and their supersymmetric vacua is interesting not just as warm-up exercise for the more complicated cases of interest in string theory but also because these theories arise in dimensional reductions of 10-dimensional superstring effective theories and the solutions found can be uplifted to full 11- or 10-dimensional M/string theory vacua still preserving a large amount of supersymmetry.

It is in this way that the supersymmetric Gödel spacetime solutions of M/string theory have been found: as a maximally supersymmetric vacuum of $N = 1, d = 5$ SUGRA which could be uplifted to 10 and 11 dimensions [2]. Many higher- [10] and also lower-dimensional

[11] examples of supersymmetric supergravity solutions with a spacetime geometry similar to the original 4-dimensional cosmological solution by K. Gödel [12] (henceforth referred to as Gödel spacetimes) have been found after it was observed that Gödel spacetimes are T-duals of H*pp*-wave spacetimes[3]. This fact has raised many questions about the consistency of string theory in such spacetimes which, being supersymmetric vacua on which (for instance) black holes can be placed [16, 17] and on which strings may be quantized [10, 18, 19], have generically closed timelike curves (CTCs). This is one of the various pathologies of General Relativity that supersymmetry or string theory are supposed to cure, forbid or explain in a consistent way and a great deal of interesting work has been done in this direction [9, 20, 21, 22, 23].

The supersymmetric vacua of M/string theory provide an interesting arena on which to study the general problems of quantization on curved spacetimes, due to the high degree of (super) symmetry they have, which, through the definition of conserved (super) charges and symmetry (super) algebra, gives a good control and understanding of the kinematics of the field theories defined on them. The fact that the aDS/CFT correspondence "works" is largely owed to the coincidence of the symmetry superalgebras of the theories involved.

The symmetry (super) algebras of the M/string theory vacua (and of the field theories defined on them) arise deformations of the symmetry (super) algebra of the original theory of which the vacua are classical solutions, which is in general the Poincaré or anti-De Sitter (super) algebra. This suggests that quantization on curved backgrounds can be, at least in some cases, equivalent to quantization using a deformation of the standard Heisenberg algebra. In other words: quantum field theories in non-trivial backgrounds can be equivalent to certain non-commutative field theories. An example of this more general relation has been provided recently by Berkovits and Seiberg [24, 8] who have found a background of Euclidean $N = 2, d = 4$ SUGRA whose symmetry superalgebra corresponds to the C-deformation introduced recently by Ooguri and Vafa [25]. We have rediscovered this background in our study of maximally supersymmetric vacua of $d = 4, 5, 6$ SUGRAs with 8 supercharges, in relation with the Gödel solution discussed above, as we are going to explain.

Our goal in this paper is to complete our knowledge of the relations between the maximally supersymmetric vacua of $d = 4, 5, 6$ SUGRAs with 8 supercharges, extending our previous work [5] to Euclidean SUGRAs which arise in timelike dimensional reductions (see Figure 1). Our main results are summarized in Figure 2 and include the identification of the above mentioned maximally supersymmetric graviphoton background of Euclidean $N = 2, d = 4$ SUGRA that cannot be obtained by Wick-rotating any known solution of the Lorentzian theory. This solution can be obtained by timelike dimensional reduction of the 5-dimensional maximally supersymmetric Gödel vacuum and has a flat Euclidean space geometry in presence of a non-trivial anti-selfdual 2-form flux which has suggested the name *flacuum* solution for it. Solutions of this kind are just as generic as those of

---

[3] Gödel spacetimes had previously been found as solutions of compactified string theory in [13, 14]. More recent solutions of supergravities in various dimensions with Gödel spacetimes have been constructed in [15].

Gödel type since they can be obtained by timelike compactifications of these and we study some examples and, in more detail, the 4-dimensional one which can also be interpreted as a $U(1)$ instanton over a T^4.

The existence of these solutions is due to the fact that the Euclidean "energy-momentum" tensor (the variation of the matter Lagrangian with respect to the metric) is not positive- or negative-definite and can vanish for non-trivial fields[4]. This vanishing can be due to a cancellation between "energy-momentum" tensors of different fields (as in the case of the Einstein-frame D-instanton solution [26, 27]) or to a cancellation between "electric" and "magnetic" components of a single field. In the case of a vector field in $d = 4$ Euclidean space, the "energy-momentum" tensor vanishes whenever the field strength is selfdual or anti-selfdual, as in the graviphoton background [24, 8].

A standard procedure to construct Euclidean theories is to dimensionally reduce a Lorentzian theory in a timelike direction. This procedure avoids the difficulties of the Wick rotation of fermions and provides a mechanism to generate flacuum solutions from higher-dimensional Lorentzian solutions with metrics of the form

$$d\hat{s}^2 = (dt + \omega)^2 - d\vec{x}_d^{\,2}\,, \tag{1.1}$$

which is precisely the general form of the supersymmetric Gödel solutions studied in [10]. This observation will allow us to construct flacuum solutions in $d = 10$ dimensions.

This paper is organized as follows: in Section 2 we summarize the dimensional reduction of standard (Lorentzian)[5] $N = (1, 0), \hat{d} = 6$ SUGRA in an arbitrary spacelike or timelike direction and its truncation to pure Lorentzian or Euclidean $N = 1, d = 5$ SUGRA. In Section 3 we summarize the dimensional reduction of the Lorentzian and Euclidean $N = 1, \hat{d} = 5$ SUGRAs obtained in the previous section in an arbitrary spacelike or timelike direction and its truncation to pure Lorentzian or Euclidean $N = 2, d = 4$ SUGRAs. The details of these reductions are explained in Appendix A, but the all the main formulae necessary for oxidation and reduction of solutions are contained in the first two sections. Observe that, as shown in Figure 1, we obtain two different Euclidean $N = 2, d = 4$ SUGRAs. These two theories are related by an analytical continuation of the vector field. In the next two sections 4 and 5 we apply the results of the first two to maximally supersymmetric vacua of $N = (1, 0), d = 6$ SUGRA to find maximally supersymmetric vacua of pure Euclidean and Lorentzian $N = 1, d = 5$ and $N = 2, d = 4$ SUGRA. We only find one not previously known from the reduction of the Gödel solution, which coincides with the graviphoton background of Seiberg and Berkovits [24, 8]. This solution, with flat space and constant anti-selfdual field strength, has interesting properties that we study in detail. In Section 6 we study more general flacuum solutions in higher dimensions. Section 8 contains our conclusions.

---

[4] The roles and definiteness of the energy and action are interchanged when we go from Lorentzian to Euclidean signature.

[5] There is no Euclidean continuation of the $N = (1, 0), \hat{d} = 6$ theory because the anti-selfduality condition of the 3-form field strength cannot be consistently defined in Euclidean signature. The minimal Euclidean SUGRA in six dimensions has not yet been constructed. It would be interesting to construct this theory since it must be related by spacelike dimensional reduction to the same Euclidean $N = 1, d = 5$ SUGRA that we are going to obtain from timelike dimensional reduction.



Figure 1: Relations between the different Euclidean (**E**) and Lorentzian (**L**) $d = 4, 5, 6$ SUGRA theories with 8 supercharges by timelike (**t**) or spacelike (**s**) dimensional reductions. There is no Euclidean continuation of the Lorentzian $N = (1, 0), d = 6$ theory, due to the anti-selfduality of its 3-form field strength, and there are two different Euclidean $N = 2, d = 4$ SUGRAs related by analytical continuation of the vector field.

2 Dimensional reduction of $N = (1, 0), d = 6$ SUGRA in the timelike direction

$N = (1, 0), d = 6$ SUGRA[6], the minimal 6-dimensional SUGRA, consists of the metric $\hat{e}^{\hat{a}}{}_{\hat{\mu}}$, 2-form field $\hat{B}^{-}_{\hat{\mu}\hat{\nu}}$ with anti-self-dual field strength $\hat{H}^{-} = 3\partial\hat{B}^{-}$ and positive-chirality symplectic Majorana-Weyl gravitino that can be described as a single, complex Dirac spinor satisfying a Weyl condition $\hat{\psi}^{+}_{\hat{\mu}}$. The bosonic equations of motion can be derived from the action

$$\hat{S} = \int d^6\hat{x}\sqrt{|\hat{g}|}\,\left[\hat{R} + \tfrac{1}{24}(\hat{H}^{-})^2\right]\,, \tag{2.1}$$

imposing afterwards the anti-self-duality constraint ${}^\star\hat{H}^{-} = -\hat{H}^{-}$. The gravitino supersymmetry transformation rule is (for zero fermions)

---

[6] Our conventions are essentially those of [5] which are those of [28] with some changes in the normalizations of the fields. In particular, we use mostly minus signature and Latin (resp. Greek or underlined) letters for Lorentz (resp. curved) indices. Further, $\hat{\gamma}_7 = \hat{\gamma}_0 \cdots \hat{\gamma}_5$, $\hat{\gamma}_7^2 = +1$, $\hat{\epsilon}^{012345} = +1$, $\hat{\gamma}^{\hat{a}_1\cdots\hat{a}_n} = \frac{(-1)^{[n/2]}}{(6-n)!}\hat{\epsilon}^{\hat{a}_1\cdots\hat{a}_n\hat{b}_1\cdots\hat{b}_{6-n}}\hat{\gamma}_{\hat{b}_1\cdots\hat{b}_{6-n}}\hat{\gamma}_7$. Positive and negative chiralities are defined by $\hat{\gamma}_7\hat{\psi}^{\pm} = \pm\hat{\psi}^{\pm}$. We use the notation $\not{F}_a \equiv \gamma^b F_{ba}$, $\not{F} \equiv \gamma^{ab}F_{ab}$ and $\not{H} \equiv \gamma^{abc}H_{abc}$.

$$\delta_{\hat{\epsilon}^+}\hat{\psi}^+_{\hat{a}} = \left(\hat{\nabla}_{\hat{a}} - \tfrac{1}{48\sqrt{2}}\,\hat{\not{H}}^-\hat{\gamma}_{\hat{a}}\right)\hat{\epsilon}^+\,. \tag{2.2}$$

We can perform the dimensional reduction[7] of $N=(1,0), \hat{d}=6$ SUGRA in one dimension can be performed simultaneously for the cases in which that dimension is timelike ($\hat{\eta}_{\sharp\sharp} \equiv \alpha_{\hat{d}} = +1$) or spacelike ($\hat{\eta}_{\sharp\sharp} \equiv \alpha_{\hat{d}} = -1$). The reduction gives $N=1, d=5$ SUGRA (the minimal 5-dimensional SUGRA) (metric $e^a{}_\mu$, *graviphoton* V_μ and symplectic-Majorana gravitino that can be described as a single complex, unconstrained Dirac spinor ψ_μ [29]) coupled to a vector multiplet consisting of a gaugino λ (the 6th component of the 6-dimensional gravitino, a real scalar (the KK one k) and a vector field W_μ. V_μ and W_μ are combinations of scalars, the KK vector field A_μ and the vector field that comes from the 6-dimensional 2-form B_μ. The identification of the right combinations is made by imposing consistency of the truncation to pure $N=1, d=5$ SUGRA[8].

The result can be stated as follows: any solution of $N=(1,0), d=6$ SUGRA satisfying the truncation constraints

$$\hat{g}_{\underline{\sharp\sharp}} = \alpha_6\,, \quad \hat{B}^-_{\mu\underline{\sharp}} = -\sqrt{2}\hat{g}_{\mu\underline{\sharp}}\,, \quad \partial_{\underline{\sharp}}\hat{\epsilon} = 0\,, \tag{2.3}$$

can be consistently reduced in the direction $x^\sharp$ to a solution of pure (Lorentzian $\alpha_6 = -1$ or Euclidean $\alpha_6 = +1$) $N=1, d=5$ SUGRA whose bosonic action and gravitino supersymmetry transformation rule are given by

$$S = \int d^5x\sqrt{|g|}\,\left[R + \tfrac{1}{4}\alpha_6 F^2 - \alpha_6\tfrac{1}{12\sqrt{3}}\tfrac{\epsilon}{\sqrt{|g|}}FFV\right]\,. \tag{2.4}$$

and

$$\delta_\epsilon\psi_a = \left\{\nabla_a - \tfrac{i}{8\sqrt{3}}\alpha_6^{3/2}(\gamma^{bc}\gamma_a + 2\gamma^b g^c{}_a)F_{bc}\right\}\epsilon\,. \tag{2.5}$$

with the same supersymmetries and with the same Killing spinor. The 5-dimensional fields are related to the 6-dimensional ones by

$$g_{\mu\nu} = \hat{g}_{\mu\nu} - \alpha_6\hat{g}_{\mu\underline{\sharp}}\hat{g}_{\nu\underline{\sharp}}\,, \qquad V_\mu = -\sqrt{3}\alpha_6\hat{g}_{\mu\underline{\sharp}}\,. \tag{2.6}$$

Conversely, any solution of Lorentzian or Euclidean $N=1, d=5$ SUGRA can be uplifted to a solution of Lorentzian $N=(1,0), \hat{d}=6$ SUGRA whose fields are given by

$$\begin{array}{rclrcl}
\hat{g}_{\underline{\sharp\sharp}} & = & \alpha_6\,, & \hat{B}_{\mu\underline{\sharp}} & = & -\sqrt{\tfrac{2}{3}}\alpha_6 V_\mu\,, \\[2ex]
\hat{g}_{\mu\underline{\sharp}} & = & \tfrac{1}{\sqrt{3}}\alpha_6 V_\mu\,, & \hat{g}_{\mu\nu} & = & g_{\mu\nu} + \tfrac{1}{3}\alpha_6 V_\mu V_\nu\,.
\end{array} \tag{2.7}$$

---

[7] In this section we use hats for 6-dimensional objects and no hats for 5-dimensional objects. In general we will use hats for higher-dimensional ($\hat{d}$-dimensional) indices and no hats for $d = \hat{d} - 1$-dimensional indices. The index corresponding to the dimension which is being reduced is $\sharp$ which will be t in the timelike case or z spacelike case. Observe that, with our mostly minus convention we get a negative-definite d-dimensional metric in the timelike case $\eta_{ab} = \hat{\eta}_{ab} = -\delta_{ab}$.

[8] The details of this dimensional reduction and truncation are explained in Appendix A.

3 Dimensional reduction of $N = 1, \hat{d} = 5$ SUGRA in the timelike direction

The dimensional reduction of $N = 1, \hat{d} = 5$ SUGRA in one dimension gives pure $N = 2, d = 4$ SUGRA (metric $e^a{}_\mu$, graviphoton V_μ and a pair gravitinos that can be combined into a complex Dirac spinor ψ_μ) coupled to a vector multiplet (a vector W_μ, a pair of scalars k and l and a complex gaugino λ) [30]. As in the previous case, the matter vector field W_μ is a combination of the scalars KK vector field and of the 5-dimensional vector that has to be identified studying the consistency of truncation in the action and supersymmetry transformation rules.

We can reduce simultaneously both the Euclidean ($\alpha_6 = +1$) and Lorentzian ($\alpha_6 = -1$) versions of $N = 1, \hat{d} = 5$ SUGRA and, further, we can do it in timelike ($\alpha_5 = +1$) and spacelike ($\alpha_5 = -1$) directions in one shot[9]. The result will be a Euclidean theory when $\alpha_6\alpha_5 = -1$ and a Lorentzian theory when $\alpha_6\alpha_5 = +1$. The reduction, identification of W_μ and consistent truncation of the vector multiplet are performed in detail in Appendix A and we obtain the following action and gravitino supersymmetry transformation rule

$$S = \int d^4x\sqrt{|g|}\left[R + \tfrac{1}{4}\alpha_6 F^2(V)\right] , \qquad \delta_\epsilon\psi_a = \left\{\nabla_a + \tfrac{i}{8}\alpha_6^{3/2} \not{F}(V)\gamma_a\right\}\epsilon\,. \tag{3.1}$$

These expressions exhibit a curious dependence on α_6 that indicates that we get two different Euclidean theories depending on the order in which we have reduced in the spacelike and timelike directions: if we compactify first in a spacelike direction $\alpha_6 = -1$ to Lorentzian $N = 1, d = 5$ SUGRA and then in a timelike direction to Euclidean $N = 2, d = 4$ SUGRA we get the same theory that we would obtain by Wick-rotating the Lorentzian theory considering the vector field V_μ as a tensor. If we compactify first in a timelike direction $\alpha_6 = +1$ to Euclidean $N = 1, d = 5$ SUGRA and then in a spacelike direction to Lorentzian $N = 2, d = 4$ SUGRA we get the same theory that we would obtain by Wick-rotating the Lorentzian theory considering V_μ as a pseudotensor. These two Euclidean theories that we denote by $N = 2_{\alpha_6}, d = 4$ are then related by an analytical continuation of the vector field $V_\mu \to iV_\mu$.

This apparent contradiction in the results of the two compactifications in one timelike and one spacelike direction is related to the well-known fact that Wick rotations and Hodge duality (used in the reduction from $d = 5$ to $d = 4$) do not commute.

The detailed calculations and results of the appendix imply that any solution of Euclidean ($\alpha_6 = +1$) or Lorentzian ($\alpha_6 = -1$) $N = 1, \hat{d} = 5$ SUGRA satisfying the truncation constraints

$$\hat{g}_{\underline{\sharp\sharp}} = \alpha_5\,, \quad \hat{V}_{\underline{\sharp}} = 0\,, \quad \sqrt{3}F(A) - {}^\star F(B) = 0\,, \quad \partial_{\underline{\sharp}}\hat{\epsilon} = 0\,, \tag{3.2}$$

---

[9] Only the case $\alpha_6 = \alpha_5 = +1$ is not possible, since we only have one time.

can be reduced, preserving all the unbroken supersymmetries, in the direction $x^{\sharp}$ to a solution of pure Euclidean or Lorentzian ($\alpha_5\alpha_6 = -1, +1$) $N = 2_{\alpha_6}, d = 4$ SUGRA with metric and vector field given by

$$g_{\mu\nu} = \hat{g}_{\mu\nu} - \alpha \hat{g}_{\mu\underline{\sharp}}\hat{g}_{\nu\underline{\sharp}}\,, \qquad V_\mu = -\tfrac{2}{\sqrt{3}}\hat{V}_\mu\,. \tag{3.3}$$

Conversely, any solution of pure Euclidean or Lorentzian $N = 2_{\alpha_6}, d = 4$ SUGRA can always be uplifted, preserving all unbroken supersymmetries, to a solution of Euclidean ($\alpha_6 = +1$) or Lorentzian ($\alpha_6 = -1$) $N = 1, \hat{d} = 5$ SUGRA whose fields are related to the 4-dimensional ones by

$$\begin{aligned}
\hat{g}_{\underline{\sharp\sharp}} &= \alpha_5\,, & \hat{V}_{\underline{\sharp}} &= 0\,,\\
\hat{g}_{\mu\underline{\sharp}} &= \alpha_5 A_\mu\,, & \hat{V}_\mu &= -\tfrac{\sqrt{3}}{2}V_\mu\,,\\
\hat{g}_{\mu\nu} &= g_{\mu\nu} + \alpha_5 A_\mu A_\nu\,,
\end{aligned} \tag{3.4}$$

where A_μ is defined in terms of V_μ by

$$F(A) = -\tfrac{1}{2}{}^{\star}F(V)\,. \tag{3.5}$$

These are the essential formulae that we have to use for oxidation and reduction of solutions and generalize those obtained in the Lorentzian cases in [5].

4 Maximally supersymmetric vacua of Euclidean $N = 1, d = 5$ SUGRA

The most interesting solutions of $N = (1,0), \hat{d} = 6$ SUGRA to which the results of the previous sections may be applied are the maximally supersymmetric vacua since their reduction should give *all* the maximally supersymmetric vacua of Euclidean and Lorentzian $N = 1, d = 5$ SUGRA. The only maximally supersymmetric vacua of $N = (1,0), \hat{d} = 6$ SUGRA are [2, 3]

1. Minkowski spacetime.
2. The 1-parameter family of Kowalski-Glikman (KG6) H*pp*-wave solutions found in Ref. [31].
3. The 1-parameter family of solutions with $aDS_3 \times S^3$ geometry found in Ref. [32] as the near-horizon limit of the self-dual string solution.

From these three solutions, all the maximally supersymmetric vacua of Lorentzian $N = 1, d = 5$ SUGRA, classified in [1] can be found by dimensional reduction in a spacelike direction [5, 2, 3, 9] (see Figure 2). They are

1. Minkowski spacetime.
2. The 1-parameter family of Kowalski-Glikman $KG5$ Hpp-wave solutions solutions found in Ref. [31].
3. The 1-parameter family of solutions with $aDS_3 \times S^2$ geometry found in Ref. [32] as near-horizon limit of the extreme string solution. It coincides with the near-horizon limit of the critically-rotating ($\jmath = 1$) BMPV black hole [33].
4. The 1-parameter family of solutions with $aDS_2 \times S^3$ geometry found in Ref. [34] as near-horizon limit of the extreme (non-rotating) black hole solution.
5. The 2-parameter family of $N = 2, d = 5$ solutions found in Ref. [35] as the near-horizon limit of the supersymmetric rotating BMPV black hole solution.
6. The 1-parameter family of Gödel-like solutions found in [1].
7. A 2-parameter family of solutions found in [1] that has just been identified as the over-rotating ($\jmath > 1$) BMPV black holes [9].

Now we want to find non-trivial maximally supersymmetric solutions of Euclidean $N = 1, d = 5$ SUGRA by dimensional reduction in a spacelike direction. As we are going to see, all of them can be obtained by Wick rotation of maximally supersymmetric vacua of Lorentzian $N = 1, d = 5$ SUGRA.

Let us consider the spacelike dimensional reductions of

4.1 The $aDS_3 \times S^3$ vacuum

aDS_3 can be seen as a spacelike $U(1)$ fibration over aDS_2 with metric (for unit radius)

$$d\Pi^2_{(3)} = \tfrac{1}{4}\left[d\Pi^2_{(2)} - (d\eta + \sinh\chi\, d\phi)^2\right] , \tag{4.1}$$

where

$$d\Pi^2_{(2)} = \cosh^2\chi d\phi^2 - d\chi^2 , \tag{4.2}$$

is the metric of aDS_2 with unit radius, just as the round S^3 can be seen as a Hopf $U(1)$ fibration over S^2 with metric (for unit radius)

$$d\Omega^2_{(3)} = \tfrac{1}{4}\left[d\Omega^2_{(2)} + (d\psi + \cos\theta d\varphi)^2\right] , \tag{4.3}$$

where

$$d\Omega^2_{(2)} = d\theta^2 + \sin^2\theta d\varphi^2 , \tag{4.4}$$

is the standard metric of S^2 with unit radius.

It is possible to perform spacelike dimensional reductions on linear combinations of these two fibers (i.e. of the coordinates η and ψ), obtaining the near-horizon limit of the extreme rotating BMPV black hole [5] (see Figure 2). The metrics of all these 5-dimensional solutions are $U(1)$ fibrations over $aDS_2 \times S^2$, the $U(1)$ fiber being orthogonal to the one we have reduced over.

It is also possible to see aDS_3 as a timelike $U(1)$ fibration over the hyperbolic plane H_2 with metric (for unit radius)

$$d\Pi^2_{(3)} = \tfrac{1}{4}\left[(dt + \cosh\chi\, d\phi)^2 - dH^2_{(2)}\right] \,, \tag{4.5}$$

where

$$dH^{(2)}_2 = \sinh^2\chi d\phi^2 + d\chi^2 \,, \tag{4.6}$$

is the standard metric of the hyperbolic plane of unit radius. We can perform further spacelike and also timelike dimensional reductions on linear combinations of these two fibers. The spacelike reductions give the near-horizon limit of the extreme over-rotating BMPV black hole [9] and here we are going to see that the timelike reductions give their analytical continuations to Euclidean signature.

The maximally supersymmetric solution with $aDS_3 \times S^3$ geometry is given by

$$\left\{\begin{array}{rcl} d\hat{s}^2 & = & R_3^2 d\Pi^2_{(3)} - R_3^2 d\Omega^2_{(3)} \,, \\ \\ \hat{H}^- & = & 4\sqrt{2} R_3 \left[\omega_{aDS_3} - \omega_{(3)}\right] , \end{array}\right. \tag{4.7}$$

where ω_{aDS_3} and $\omega_{(3)}$ are the volume forms of the unit radius aDS_3 and S^3 metrics written above. Using the timelike $U(1)$ fibration Eq. (4.5) we find that a convenient gauge for the 2-form potential is

$$\hat{B}^- = -R_3/\sqrt{2}\left[\cosh\chi d\phi \wedge dt - \cos\theta d\varphi \wedge d\psi\right] \,. \tag{4.8}$$

Rescaling t and ψ by $2/R_3$ and boosting the metric in the direction ψ

$$\begin{array}{rcl} t & = & 2/R_3(\cosh\xi\, t' + \sinh\xi\, \psi') \,, \\ \\ \psi & = & 2/R_3(\sinh\xi\, t' + \cosh\xi\, \psi') \,, \end{array} \tag{4.9}$$

the solution takes the form

$$\left\{\begin{array}{rcl} d\hat{s}^2 & = & (dt' + A^{(t)})^2 - (d\psi' + A^{(\psi)})^2 - (R_3/2)^2[dH^2_{(2)} + d\Omega^2_{(2)}] \,, \\ \\ \hat{B}^- & = & -\sqrt{2} A^{(t)} \wedge dt' + \sqrt{2} A^{(\psi)} \wedge d\psi' \,, \end{array}\right. \tag{4.10}$$

where $A^{(t,\psi)}$ are the 1-forms

$$
\begin{aligned}
A^{(t)} &= R_3/2(\cosh\xi\cosh\chi d\phi - \sinh\xi\cos\theta d\varphi)\,,\\
A^{(\psi)} &= R_3/2(\cosh\xi\cos\theta d\varphi - \sinh\xi\cosh\chi d\phi)\,.
\end{aligned}
\tag{4.11}
$$

The truncation constraints Eqs. (2.3) are satisfied for $x^\sharp = t'$ and $x^\sharp = \psi'$ and, thus, we can dimensionally reduce this solution to a maximally supersymmetric solution of Euclidean or Lorentzian $N = 1, d = 5$ SUGRA. In the spacelike case we get

$$
\left\{
\begin{aligned}
ds^2 &= (dt' + A^{(t)})^2 - (R_3/2)^2[dH^2_{(2)} + d\Omega^2_{(2)}]\,,\\
V &= -\sqrt{3}A^{(\psi)}\,, \qquad F = -\tfrac{\sqrt{3}}{2}R_3(\sinh\xi\omega_{aDS_2} + \cosh\xi\omega_{(2)})\,,
\end{aligned}
\right.
\tag{4.12}
$$

which is the near-horizon limit of the extreme critically rotating and over-rotating BMPV black hole [9]. $\jmath = \cosh\xi$ is the rotation parameter. The critically rotating case has $\jmath = 1$, i.e. $\sinh\xi = 0$ and the fibration takes place only over the hyperbolic plane, giving aDS_3. The total metric describes the solution $aDS_3 \times S_2$. For any other value of ξ, $\jmath > 1$, and we are in the over-rotating case.

In the timelike case, we get

$$
\left\{
\begin{aligned}
ds^2 &= -(d\psi' + A^{(\psi)})^2 - (R_3/2)^2[dH^2_{(2)} + d\Omega^2_{(2)}]\,,\\
V &= -\sqrt{3}A^{(t)}\,, \qquad F = -\tfrac{\sqrt{3}}{2}R_3(\cosh\xi\omega_{H_2} + \sinh\xi\omega_{(2)})\,.
\end{aligned}
\right.
\tag{4.13}
$$

These two solutions are related by a Wick rotation $t \to i\psi$ $V \to iV$ which has to be accompanied of $\xi \to \xi - i\pi/2$ to make V real again. The Euclidean solution can also be obtained by a Wick rotation $t \to i\psi$ $V \to iV$ from the near-horizon limit of the extreme BMPV black hole accompanied of $\xi \to i\xi$ to make V real again.

On the other hand, these solutions can be seen, respectively, as a timelike $\mathbb{R}$ or spacelike $U(1)$ fibration over $H_2 \times S^2$.

Let us now consider the timelike dimensional reductions of

4.2 The $KG6$ Hpp-wave vacuum

As shown in [5], the KG6 solution can be conveniently written substituting g_{uu} completely by a Sagnac connection 1-form $\omega = \lambda(x^1dx^2 - x^3dx^4)$ whose main property is that $d\omega$ is anti-selfdual in the 4-dimensional Euclidean space spanned by $\vec{x}_4 = (x^1, x^2, x^3, x^4)$:

$$
\left\{
\begin{aligned}
d\hat{s}^2 &= 2du(dv + \sqrt{2}\omega) - d\vec{x}^2_4\,,\\
\hat{B}^- &= -2\omega \wedge du\,,
\end{aligned}
\right.
\tag{4.14}
$$

and, using $u = (t + z)/sqrt2$, we can rewrite it in the form

$$
\left\{
\begin{array}{rcl}
d\hat{s}^2 & = & (dt+\omega)^2 - (dz-\omega)^2 - d\vec{x}_4^2\,, \\
 & & \\
\hat{B}^- & = & -\sqrt{2}\omega\wedge dt - \sqrt{2}\omega\wedge dz\,,
\end{array}
\right.
\tag{4.15}
$$

that shows that the truncation constraints Eqs. (2.3) are satisfied for $x^\sharp = t$ and $x^\sharp = z$ and, thus, we can dimensionally reduce this solution to maximally supersymmetric solutions of Euclidean or Lorentzian $N=1, d=5$ SUGRA. The spacelike dimensional reduction gives the Gödel solution

$$
\left\{
\begin{array}{rcl}
ds^2 & = & (dt+\omega)^2 - d\vec{x}_4^2\,, \\
 & & \\
V & = & -\sqrt{3}\omega\,,
\end{array}
\right.
\tag{4.16}
$$

and the timelike reduction gives its Euclidean version

$$
\left\{
\begin{array}{rcl}
ds^2 & = & -(dz-\omega)^2 - d\vec{x}_4^2\,, \\
 & & \\
V & = & -\sqrt{3}\omega\,,
\end{array}
\right.
\tag{4.17}
$$

that can be obtained by Wick-rotating $t \to iz$, $V \to iV$ plus $\lambda \to -i\lambda$ to make V real. These two solutions can also be seen as timelike $\mathbb{R}$ or spacelike $U(1)$ fibrations over the 4-dimensional flat Euclidean space.

No more timelike dimensional reductions satisfying the truncation constraints seem to be possible, and, thus, we do not expect any more maximally supersymmetric solutions of Euclidean $N=1, d=5$ SUGRA.

5 Maximally supersymmetric vacua of Euclidean $N = 2_\pm, d=4$ SUGRA

Next, we are going to dimensionally reduce the maximally supersymmetric vacua of Euclidean and Lorentzian $N=1, d=5$ SUGRA to find maximally supersymmetric vacua of Euclidean $N=2_\pm, d=4$ SUGRAs. The maximally supersymmetric vacua of Lorentzian $N=2, d=4$ SUGRA[10] have been obtained by spacelike dimensional reduction of the vacua of Lorentzian $N=2, d=5$ SUGRA listed in Section 4 [5].

The only Lorentzian solutions that satisfy the truncation constraints for timelike reduction seem to be Minkowski spacetime, the near-horizon (NH) limit of the critically rotating and over-rotating BMPV black holes (BH) Eq. (4.12) and the Gödel solution Eq. (4.16).

---

[10] All the solutions of Lorentzian $N=2, d=4$ SUGRA that preserve some supersymmetry were found in [36]. The maximally supersymmetric vacua of the theory are just three: Minkowski spacetime, the dyonic Robinson-Bertotti solution [7] and the KG4 H*pp*-wave solution [6].

The only Euclidean solutions that satisfy the analogous truncation constraints are the Euclidean NH limit of the rotating BMPV BH[11] Eq. (4.13) and the Euclidean Gödel solution Eq. (4.17) Let us study the non-trivial cases.

5.1 NH limits of the extreme critically rotating and over-rotating BMPV BHs

Using the formulae of Section 3 on the solution Eq. (4.12) we immediately get a maximally supersymmetric solution of Euclidean $N = 2_{-}, d = 4$ SUGRA

$$\left\{\begin{array}{rcl} ds^2 & = & -(R_3/2)^2[dH_2^2 + d\Omega_2^2]\,, \\ \\ F & = & R_3\left(\sinh\xi\omega_{H_2} + \cosh\xi\omega_{(2)}\right)\,, \end{array}\right. \tag{5.1}$$

which is the Wick-rotated ($\phi = -it$) version of the well-known dyonic Robinson-Bertotti solution [7]. Observe that, in order to have a real Euclidean solution, the Lorentzian electric-magnetic rotation angle has also been Wick-rotated $\xi \to i\xi$.

5.2 The Euclidean NH limits of extreme rotating BMPV BHs

Using the formulae of Section 3 on the solution Eq. (4.13) we immediately get a maximally supersymmetric solution of Euclidean $N = 2_{+}, d = 4$ SUGRA

$$\left\{\begin{array}{rcl} ds^2 & = & -(R_3/2)^2[dH_2^2 + d\Omega_2^2]\,, \\ \\ F & = & R_3\left(\cosh\xi\omega_{H_2} + \sinh\xi\omega_{(2)}\right)\,, \end{array}\right. \tag{5.2}$$

This solution is related to the Euclidean dyonic Robinson-Bertotti solution by analytical continuation of the vector field $V \to iV$ together with $\xi \to \xi - i\pi/2$.

5.3 The Euclidean and Lorentzian Gödel solutions and the flacuum solution

The $\hat{d} = 5$ Gödel solutions are given in Eqs. (4.16) and (4.17). The formulae of Section 3 give, in both cases, the remarkable maximally supersymmetric 4-dimensional vacuum solution with constant anti-selfdual flux (*flacuum*) found in [8]

$$\left\{\begin{array}{rcl} ds^2 & = & -d\vec{x}_4^{\,2}\,, \\ \\ V & = & 2\omega\,. \end{array}\right. \tag{5.3}$$

The triviality of the metric in presence of non-trivial matter fields is surprising at first sight, but it is not the first example of non-trivial Euclidean SUGRA solution with flat spacetime: the Einstein-frame D-instanton metric [27] (and that of its 4-dimensional analogue [26]) is also flat. In all these cases, the Euclidean version of the energy-momentum tensor vanishes identically, but for different reasons. In the D-instanton case the dilaton and RR 0-form energy-momentum tensors cancel each other on-shell. In the present case, anti-selfduality of the vector field strength makes the energy-momentum tensor vanish identically. This can be easily seen using Eq. (A.28) which tells us that, actually, the Euclidean version of energy-momentum tensor for any (anti-) selfdual Abelian or non-Abelian vector field vanishes identically and, thus, decouples from the metric[12]. This implies that solutions such as the BPST instanton [37] are not just solutions of the Euclidean Yang-Mills equations on S^4 with the standard metric, but they are also solutions of the full supergravity equations, provided that the metric satisfies the vacuum Einstein equations. We will discuss this and similar solutions in the next section.

---

[11] As we have seen, the Wick-rotated NH limits of the extreme over-rotating and un "under-rotating" BMPV BHs are identical.

The above solution is guaranteed to be maximally supersymmetric in both Euclidean $N = 2_{\pm}, d = 4$ SUGRA. It is worth studying how the Killing spinors integrability conditions are satisfied. They take the form[13]

$$\left\{-\tfrac{1}{4}R_{\mu\nu}{}^{ab}\gamma_{ab} + \tfrac{1}{4}T_{[\mu|\rho|}\gamma^{\rho}{}_{\nu]} + \tfrac{1}{4}\ \not{\nabla}\left(F_{\mu\nu} + {}^{\star}F_{\mu\nu}\gamma^{\sharp}\right)\right\}\epsilon = 0\,. \tag{5.4}$$

In the case at hand the first term vanishes identically because the space is flat, the second because of the anti-selfduality of the vector field strength and the third and fourth because the field strength is covariantly constant (constant in Cartesian coordinates).

(Anti-) selfduality implies flatness and, then, covariant constancy determines a unique solution (up to $SO(4)$ rotations). Suppressing this last requirement allows for more solutions[14] which would only preserve a half of the supersymmetries.

This solution preserves all supersymmetries and the Killing spinors depend on all coordinates:

$$\epsilon(x) = \exp\{-\tfrac{1}{8}\ \not{F}\not{x}\}\epsilon_0\,, \tag{5.5}$$

where ϵ_0 is a constant spinor and $\not{x} \equiv x^{\mu}\gamma_{\mu}$. Using the anti-selfduality of F, we can rewrite the exponent in this form

$$-\tfrac{1}{8}\ \not{F}\not{x} = -\tfrac{1}{4}\ \not{F}_{\mu}x^{\mu}(1-\gamma^{\sharp})\,, \tag{5.6}$$

where $(\gamma^{\sharp})^2 = 1$ in this case. This implies that $(\not{F}\not{x})^2 = 0$ and, thus, the Killing spinors can be written in the form

$$\epsilon(x) = \{1 - \tfrac{1}{4}\ \not{F}_{\mu}x^{\mu}(1-\gamma^{\sharp})\}\epsilon_0\,. \tag{5.7}$$

If we split our complex Dirac spinors into chiral halves, we see that the positive chirality Killing spinor is simply constant, as in empty Euclidean space, and the negative chirality Killing spinor has a x-dependent deformation with respect to that of empty Euclidean space which induces a deformation of the supersymmetry algebra [24, 8] that affects only to negative-chirality objects. Using the methods of [38] and the notation of [39, 40], and using 4-component complex (Dirac) spinors, we find the following non-vanishing (anti-) commutators

---

[12] But not the other way around since (anti-) selfduality is established with respect to a given metric.

[13] For simplicity we consider the $\alpha_6 = -1$ case.

[14] For instance, $A = dx^{\sharp}/r + \cos\theta d\varphi$ in spherical coordinates.

$$\begin{aligned}
\left\{\mathcal{Q}^{\dagger}_{(\alpha)}, \mathcal{Q}_{(\beta)}\right\} &= (\gamma^1\gamma^a)_{\alpha\beta}P_{(a)} - [\gamma^1\tfrac{1}{2}(1-\gamma^{\sharp})]_{\alpha\beta}M\,, \\
\left[\mathcal{Q}_{(\alpha)}, P_{(a)}\right] &= -\mathcal{Q}_{(\beta)}\Gamma_s(P_{(a)})^{\beta}{}_{\alpha}\,, \\
\left[\mathcal{Q}_{(\alpha)}, M\right] &= -\mathcal{Q}_{(\beta)}\Gamma_s(M)^{\beta}{}_{\alpha}\,, \\
\left[P_{(a)}, M\right] &= -P_{(b)}\Gamma_v(M)^{b}{}_{a}\,.
\end{aligned} \tag{5.8}$$

The bosonic generators $P_{(a)}$ of this algebra are associated to the translational Killing vectors ∂_a. The bosonic generator M is a Lorentz ($SO(4)$) rotation $\frac{1}{2}F^{ab}M_{ab}$ where the Lorentz generators M_{ab} are associated the Killing vectors $-2x_{[a}\partial_{b]}$. Both the P_as and M act on spinors through their spinorial representations

$$\Gamma_s(P_{(a)})^{\beta}{}_{\alpha} = \left[\tfrac{1}{4}\not{F}_a(1-\gamma^{\sharp})\right]^{\beta}{}_{\alpha}\,, \qquad \Gamma_s(M)^{\beta}{}_{\alpha} = \left[\tfrac{1}{2}\not{F}\right]^{\beta}{}_{\alpha} = \left[\tfrac{1}{4}\not{F}(1+\gamma^{\sharp}\right]^{\beta}{}_{\alpha}\,, \tag{5.9}$$

and on vectors through the vector representation

$$\Gamma_v(M)^{b}{}_{a} = -F^{b}{}_{a}\,. \tag{5.10}$$

This superalgebra is a deformation of the standard Poincaré $N=2, d=4$ superalgebra. This is a common property of all the maximally supersymmetric vacua of $N=2, d=4$ SUGRA. However, this deformation is particularly interesting because the bosonic translations algebra is not modified and, actually, only the anticommutator of the negative chirality part of the supercharge is deformed and this deformation induces a deformation of the anticommutator of the fermionic superspace coordinates of negative chirality [24, 8].

The relation between the flacuum and Gödel solutions (whose Killing spinors are, according to our results, absolutely identical) implies a relation between their superalgebras which is important to understand. The only difference between them is the occurrence in the Gödel superalgebra of the bosonic generator $P_{(0)}$ associated to time translations in the anticommutator of the supercharges and, more importantly, in the commutator of the generators associated to space translations $P_{(a)}$. If we write the Gödel solution Eq. (4.16) in the gauge in which the 1-form ω takes the form

$$\omega = \tfrac{1}{2}F_{ab}x^a dx^b\,, \quad a,b = 1,2,3,4\,, \quad F_{12} = -F_{34} = \lambda\,, \tag{5.11}$$

we find the Killing vectors[15]

---

[15] There are more Killing vectors, but they are not relevant insofar they do not appear in the anticommutator of the supercharges.

$$\partial_t\,, \qquad \partial_a - \tfrac{1}{2}F_{ab}x^b\partial_t\,, \qquad -\tfrac{1}{2}F^{ab}x_a\partial_b\,, \tag{5.12}$$

associated, respectively, to the generators of time and space translations $P_{(0)}$ and $P_{(a)}$ and to the rotation M. The non-vanishing (anti-) commutators of the Gödel superalgebra are

$$\begin{aligned}
\left\{\mathcal{Q}^\dagger_{(\alpha)}, \mathcal{Q}_{(\beta)}\right\} &= (\gamma^1\gamma^a)_{\alpha\beta}P_{(a)} + (\gamma^1\gamma^\sharp)_{\alpha\beta}P_{(0)} - [\gamma^1\tfrac{1}{2}(1-\gamma^\sharp)]_{\alpha\beta}M\,,\\
\left[\mathcal{Q}_{(\alpha)}, P_{(a)}\right] &= -\mathcal{Q}_{(\beta)}\Gamma_s(P_{(a)})^\beta{}_\alpha\,,\\
\left[\mathcal{Q}_{(\alpha)}, M\right] &= -\mathcal{Q}_{(\beta)}\Gamma_s(M)^\beta{}_\alpha\,,\\
\left[P_{(a)}, M\right] &= -P_{(b)}\Gamma_v(M)^b{}_a\,, \qquad \left[P_{(a)}, P_{(b)}\right] = F_{ab}P_{(0)}\,.
\end{aligned} \tag{5.13}$$

In this background space translations commute only up to a time translation, because space translations δx^a have to be compensated by a time translation $\delta t = \frac{1}{2}F^{ab}x_b$ to leave the metric invariant. This deformation is interesting because is suggests that string theory quantized in the Gödel background will give a non-commutative theory in which, instead of modifying the commutator of positions, it is the commutator of momenta which are deformed as above. Supersymmetry requires a corresponding modification of the anticommutator f the supercharges as in the flacuum solution.

In the dimensional reduction to the flacuum solution the generator $P_{(0)}$ becomes a central charge that generates gauge transformations of the Kaluza-Klein vector field. Actually, it is easy to check that space translations only leave the vector field of the flacuum solution up to gauge transformations with gauge parameter $\frac{1}{2}F^{ab}x_b$. In this sense, the flacuum superalgebra is not complete and one should add $P_{(0)}$s as in the Gödel superalgebra. However, if we are considering only t-independent 5-dimensional configurations which correspond to solutions with zero $P_{(0)}$ charge, the difference is immaterial. In $N=2, d=4$ SUGRA there are no charged perturbative states in $N=2, d=4$ SUGRA, but non-perturbative states that would be associated to 5-dimensional KK modes with non-trivial time dependence are charged with respect to $P_{(0)}$ and would feel the non-commutativity of the momenta.

5.4 The flacuum solution as an Abelian instanton

Another interesting aspect of the flacuum solution is its possible interpretation as an instanton. This is only possible if the space is compactified into a T^4 to make the action integral finite. This is, also, a base space on which the $U(1)$ gauge group can be non-trivially fibered. Euclidean Yang-Mills solutions on T^4 have been thoroughly studied in the literature[16] and we are simply going to apply the known results to the present case.

---

[16] See, for instance the pedagogical review [41] whose notation we will loosely follow. In particular we use the period vectors $(\hat{a})^c \equiv \delta_{(a)}{}^c l^{(a)}$, where summation is not to be made over indices within parenthesis.

To compactify the solution on T^4 we take the quotient of $\mathbb{R}^4$ by the $\mathbb{Z}^4$ Abelian group of discrete translations along the four coordinates x^a with periods l^a. The vector field of our solution, which we rewrite for convenience in a new gauge

$$V = \lambda(x^1 dx^2 - x^2 dx^1 - x^3 dx^4 + x^4 dx^3) \equiv F_{ab} x^a dx^b \,, \tag{5.14}$$

is not strictly periodic on T^4: when we move around the a-th period from x to $x + \hat{a}$ it changes by a gauge transformation

$$V(x + \hat{a}) = V(x) + d\Lambda_a(x) \,, \qquad \Lambda_a(x) = l^{(a)} F_{(a)b} x^b \,, \tag{5.15}$$

where $\Lambda_a(x)$ are the $U(1)$ parameters, defined modulo 2π. Consistency requires that $V(x + \hat{a} + \hat{b}) = V(x + \hat{b} + \hat{a})$, that is

$$\Lambda_a(x + \hat{b}) + \Lambda_b(x) = \Lambda_b(x + \hat{a}) + \Lambda_a(x) \mod(2\pi) \,, \tag{5.16}$$

which in our case implies

$$\lambda l^1 l^2 = \pi n \,, \qquad \lambda l^3 l^4 = \pi m \,, \tag{5.17}$$

for two integers n, m that label the possible non-trivial bundles, the topological number being given by their product nm. The Euclidean action of these SUGRA solutions is, therefore, in our conventions

$$S = -4\pi^2 |nm| \,. \tag{5.18}$$

Let us consider now taking the quotient on the spinor bundle, which requires making the identifications

$$\epsilon(x + \hat{a}) = \exp\{-\tfrac{l^{(a)}}{8} \not{F} \gamma_{(a)}\} \epsilon(x) \sim \epsilon(x) \,, \tag{5.19}$$

consistently. These transformations are not $SO(4)$ rotations but are still elements of the bigger holonomy group of the supercovariant derivative of $N = 2, d = 4$ SUGRA, which, by arguments similar to those in [42, 43] can be shown to be $SL(4, \mathbb{R})$, and, in this sense, from this more general point of view, they should be admissible.

On the other hand, the consistency of these identifications is ensured by the fact that the above transformations are indeed a representation of the commutative $\mathbb{Z}_4$ group of discrete translations: the generators of translations in the x^a directions, $P_{(a)}$ are represented on the spinors by the mutually commuting operators $\Gamma_s(P_{(a)})$ given in Eq. (5.9).

The consistency of this construction has interesting implications: the compactness of the gauge group ($U(1)$ instead of $\mathbb{R}$), reflected by the periodicity of the gauge parameters $\Lambda_a(x)$ implies the compactness of the $\hat{d} = 5$ time coordinate. All the vacua of $N = 1, \hat{d} = 5$ SUGRA (except the KG5 H*pp*-wave) seem to be non-trivial spacelike or timelike fibrations over 4-dimensional Euclidean space or over $H_2 \times S^2$.

In this respect, it is remarkable that the only solutions that can be reduced consistently in the timelike direction (preserving all supersymmetries) have closed timelike curves or (in the $\jmath = 1$ closed lightlike curves.



Figure 2: Relations between the $d = 4, 5, 6$ vacua with 8 supercharges. Dimensional reduction in spacelike direction is represented by arrows with black arrowheads and in timelike directions with white arrowheads. The ellipse and hyperbolas represent the families of near-horizon limits of rotating and over-rotating extreme BMPV black holes. In particular, the elements α and β of these families have the geometries of $aDS_2 \times S^3$ and $aDS_3 \times S^2$. Dotted lines indicate that the corresponding solutions are related by a Penrose-Güven-like limiting procedure.

6 (Anti-) selfdual vacua in Euclidean SUGRA

It is not difficult to construct generalizations of the flacuum solution. Let us consider a $(p+1)$ form potential $A_{(p+1)}$ with field strength $F_{(p+2)} = dA_{(p+1)}$ coupled to d-dimensional gravity:

$$S = \int d^dx \sqrt{|g|}\left[R - \tfrac{(-1)^p}{2\cdot(p+2)!}F^2_{(p+2)}\right] . \tag{6.1}$$

The Einstein equation is

$$R_{\mu\nu} - \tfrac{1}{2}g_{\mu\nu}R = \tfrac{(-1)^p}{2\cdot(p+1)!}T^{(p+1)}_{\mu\nu} , \tag{6.2}$$

where the energy-momentum tensor is given by

$$T^{(p+1)}_{\mu\nu} = F_{(p+2)\,\mu}{}^{\rho_1\cdots\rho_{p+1}}F_{(p+2)\,\nu\rho_1\cdots\rho_{p+1}} - \tfrac{1}{2\cdot(p+2)}g_{\mu\nu}F^2_{(p+2)} , \tag{6.3}$$

or, equivalently, by the more useful expression

$$T^{(p+1)}_{\mu\nu} = \tfrac{1}{2}\left[F_{(p+2)\,\mu}{}^{\rho_1\cdots\rho_{p+1}}F_{(p+2)\,\nu\rho_1\cdots\rho_{p+1}} - \alpha\,{}^{\star}F_{(p+2)\,\mu}{}^{\rho_1\cdots\rho_{\tilde{p}+1}}\,{}^{\star}F_{(p+2)\,\nu\rho_1\cdots\rho_{\tilde{p}+1}}\right] , \tag{6.4}$$

where $\tilde{p} = d - p - 4$.

We are interested in solutions with vanishing energy-momentum tensor. A necessary condition for the energy-momentum tensor to vanish is that its trace does. Using (6.3) we find that

$$T_{\mu}^{(p+1)\mu} \sim \left[1 - \tfrac{d}{2(p+2)}\right] F_{(p+2)}^2 \,, \tag{6.5}$$

which vanishes when $F_{(p+2)}^2 = 0$ or when $p + 2 = d/2$. For non-trivial potentials, the first case is only possible in Lorentzian signature, but we know that the Lorentzian energy-momentum tensor only vanishes for vanishing field strengths. Thus, a vanishing energy-momentum tensor is only possible in Euclidean signature for $d = 2(p+2)$ and, using (6.4), it requires

$$F_{(p+2)\,\mu}{}^{\rho_1\cdots\rho_{p+1}} F_{(p+2)\,\nu\rho_1\cdots\rho_{p+1}} = {}^{\star}F_{(p+2)\,\mu}{}^{\rho_1\cdots\rho_{p+1}\star} F_{(p+2)\,\nu\rho_1\cdots\rho_{p+1}} \,, \tag{6.6}$$

which can be solved in any even dimension: in $d = 4n$ using (anti-) selfdual field strengths (as we have seen in $d = 4$) and in $d = 4n + 2$ using configurations which would be (anti-) self-dual in Lorentzian signature. For example, in $d = 6$ we could use $F_{(3)\,123} = F_{(3)\,456} = \lambda$ which has $F_{(3)\,123} = -{}^{\star}F_{(3)\,123}$ but $F_{(3)\,123} = +{}^{\star}F_{(3)\,123}$ (i.e. not definite selfduality properties) but clearly satisfies the above condition.

Unfortunately, these simple *Ansätze* for flacuum solutions do not work in presence of scalars such as the string dilaton that couple to $F_{(p+2)}^2$. Still, since the sign of the Euclidean "kinetic" terms and energy-momentum tensors is not uniquely defined[17], it is possible that the energy-momentum tensors and their contributions to the source for the dilaton of two fields cancel each other. Examples of these solutions are provided by the dimensional reduction in the time direction of the supersymmetric Gödel solutions of $N = 1, d = 11$ SUGRA found in [10]. These solutions take the form

$$\left\{ \begin{array}{rcl} d\hat{s}^2 & = & (dt + c_i \omega^i)^2 - d\vec{x}_{10}^{\,2} \,, \\ \\ \hat{G} & = & \tfrac{1}{4} a_{ij} d\omega^i \wedge d\omega^j \,, \end{array} \right. \tag{6.7}$$

where the ω^is are 1-forms defined by

$$\omega^i = x^{(i)} dx^{5+(i)} - x^{5+(i)} dx^{(i)} \,, \tag{6.8}$$

and where the constants c_i and a_{ij} (with $a_{ii} = 0$) have to satisfy the following two conditions:

$$a_{(i)j} a_{j(i)} = 4c_i^2 + c_j c_j \,, \qquad a_{ij} c_j = -\tfrac{1}{8} \eta^{ijklm} a_{jk} a_{lm} \,, \tag{6.9}$$

where the η^{ijklm} symbol is one is all its indices take different values and zero otherwise. It is convenient to label the solutions of this family by the two vectors of constants

---

[17] They depend on whether the Lorentzian field is a tensor or a pseudotensor, as the KK vector generically turns out to be.

$$(c_1, c_2, c_3, c_4, c_5)\,, \qquad (a_{12}, a_{13}, a_{14}, a_{15}, a_{23}, a_{24}, a_{25}, a_{34}, a_{35}, a_{45})\,. \tag{6.10}$$

The supersymmetry of some explicit Gödel solutions was studied in [10] and some examples are given in the table 1

	(c_i)	(a_{ij})	# Susies
(i)	$\lambda(1,1,0,0,0)$	$\lambda(0,-2,-2,-2,-2,-2,-2,0,0,0)$	20
(ii)	$\lambda(1,1,1,1,0)$	$\lambda(4,0,0,-2,0,0,-2,4,-2,-2,-2)$	0
(iii)	$\lambda(1,1,1,-1,0)$	$\lambda(4,0,0,-2,0,0,-2,4,-2,-2,+2)$	20
(iv)	$\lambda(1,0,0,0,0)$	$\lambda(2,0,0,-2,0,0,0,2,0,0)$	8
(v)	$\lambda(2,1,1,1,1)$	$\lambda(-6,2,2,2,0,0,0,4,4,4)$	18

Table 1: Some Gödel solutions of $N=1, d=11$ SUGRA. The first example gives, after dimensional reduction, the maximally supersymmetric Gödel solution of $N=1, d=5$ SUGRA Eq. (4.16) and, therefore, it also gives rise to the flacuum solution Eq. (5.3).

It is clear that the timelike reduction of any of these Gödel spacetimes gives a flacuum solutions of Euclidean type IIA SUGRA with flat metric and RR 4-form and 1-form given by

$$\left\{ \begin{array}{rcl} ds^2 & = & -d\vec{x}^{\,2}_{10}\,, \\ & & \\ G^{(2)} & = & c_i d\omega^i\,, \\ & & \\ G^{(4)} & = & \frac{1}{4} a_{ij} d\omega^i \wedge d\omega^j\,. \end{array} \right. \tag{6.11}$$

It is important to understand how the vanishing of the energy-momentum tensor takes place and why the dilaton is constant in this case. The action of the Euclidean type II SUGRA obtained by dimensional reduction from standard, Lorentzian 11-dimensional SUGRA is given by

$$\begin{array}{rcl} S & = & \displaystyle\int d^{10}x\sqrt{|g|}\Big\{ e^{-2\phi}\left[R - 4\left(\partial\phi\right)^2 - \tfrac{1}{2\cdot 3!}\alpha_{11}H^2\right] \;+\; \tfrac{1}{4}\alpha_{11}G^{(2)\,2} \;-\; \tfrac{1}{2\cdot 4!}G^{(4)\,2} \\ & & \qquad\qquad -\tfrac{1}{144}\tfrac{1}{\sqrt{|g|}}\,\epsilon\partial C^{(3)}\partial C^{(3)}B\Big\} \end{array} \tag{6.12}$$

This action differs from the type IIA action by the occurrences of α_{11}'s, the definition of the various field-strengths staying the same[18]. Observe that, to obtain this bosonic action by a standard Wick rotation one has to consider both the RR 1-form and the NSNS 2-form pseudotensors.

The RR 1- and 3-form kinetic terms $G^{(2)\,2}$ and $G^{(4)\,2}$ have opposite signs when $\alpha_{11}=+1$ and the dilaton does not couple directly to them in the string frame, but only through the

---

[18] Our conventions here are the same as in Ref. [44].

Ricci scalar. Thus, the exact cancellation between the respective "energy-momentum" tensors which makes the metric flat is enough to render the dilaton constant.

To end this section, let us observe that the BPST instanton [37] also provides an example of flacuum solution since its energy-momentum tensor is identically zero because of the (anti-) selfduality of its field strength on the round S^4. Thus, it is a solution of the Euclidean Einstein-Yang-Mills theory with positive cosmological constant proportional to the square inverse radius of the sphere. The embedding of this solution in a supergravity theory is more problematic because a DS-type gauged SUGRA would have to be used. It should be possible to uplift the solution to Euclidean purely gravitational $d = 7$ solution with the S^7 metric [45]

6.1 *pp*-waves, lightlike reductions and flacuum solutions

The fact that the $N = 2, d = 4$ flacuum solution originates in a timelike and spacelike dimensional reduction of the KG6 H*pp* wave suggests that there must exist a relation between this kind of solutions and *pp*-waves.

pp-wave metrics can always be put in the form

$$d\hat{s}^2 = 2du(dv + Kdu + A_{\underline{i}}dx^i) + g_{\underline{ij}}dx^i dx^j \,. \tag{6.13}$$

where all the functions in the metric are independent of v. K or $A_{\underline{i}}$ can be removed by coordinate transformations that preserve the above form of the metric, but we will keep both. Furthermore, we are going to assume independence on the second light-cone coordinate u

The components of the $\hat{d}$-dimensional Ricci tensor are

$$\hat{R}_{\underline{ij}} = R_{\underline{ij}} \,, \qquad 2\hat{R}_{\underline{iu}} = \nabla^{\underline{j}} F_{\underline{ji}} \,, \qquad \hat{R}_{\underline{uu}} = \nabla_{\underline{i}}\partial^{\underline{i}} K - \tfrac{1}{4}F^2 \,, \tag{6.14}$$

where all the objects without hats are calculated with the $d = (\hat{d} - 2)$-dimensional Euclidean wavefront metric $g_{\underline{ij}}$ and where $F_{\underline{ij}} = 2\partial_{[\underline{i}}A_{\underline{j}]}$. The Ricci scalar is just $\hat{R} = R$ and, therefore, $\hat{d}$-dimensional vacuum Einstein equations $\hat{G}_{\hat{\mu}\hat{\nu}} = 0$ imply the d-dimensional vacuum Einstein equations $G_{\underline{ij}} = 0$, the vacuum Maxwell equation in the background metric $\nabla_{\underline{j}} F^{\underline{ji}} = 0$ and the scalar equation of motion $\nabla^2 K = \frac{1}{4}F^2$.

This is similar to the result of a Kaluza-Klein reduction, except for the fact that neither the scalar nor the vector field contribute to the Euclidean "energy-momentum" tensor[19] as in flacuum solutions. Actually, the flacuum solution can be embedded in a purely gravitational $\hat{d} = 6$ *pp*-wave with $K \sim \lambda|\vec{x}_4|^2$. Another wide class of similar solutions is given by

$$F_{\underline{i\sharp}} = 2\nabla_{\underline{i}} H \,, \qquad F_{\underline{ij}} = \epsilon_{ijk}\frac{1}{\sqrt{|g|}}\nabla_{\underline{k}} H \qquad K = H^2 \,, \qquad \nabla^2 H = 0 \,. \tag{6.15}$$

---

[19] Since either of them can be removed by reparametrizations, only one of them may contribute.

It is possible to perform dimensional reductions in one lightlike dimension [46], but it does not very useful as the resulting theories have singular metric. The above observation suggests that lightlike reductions should be made in the two light-cone coordinates to give a Euclidean $\hat{d}-2$-dimensional theory that has essentially a metric an a vector field constrained to have vanishing energy-momentum tensor and to satisfy the constraint $F^2 = 4\nabla^2 K$ for some scalar K. In $d = 4$ it is possible to define Euclidean theories in which the selfdual or anti-selfdual parts of the spin connection are the only variables. It might be possible to construct a theory in which only the selfdual or anti-selfdual part of F occurs and we can speculate that this kind of theory is the one associated to lightlike dimensional reductions.

7 Limiting Solutions

Apart from the dimensional reduction from the KG6 solution, there is another way of obtaining the Gödel solution through a limiting procedure which is very similar to the Penrose-Güven limit [47, 48] that relates $aDS_n \times S^m$ supergravity vacua to Kowalski-Glikman-type Hpp-wave solutions and in general any supergravity solution to a plane wave solution with equal or larger number of isometries and supersymmetries [49, 50].

Let us consider the NH limit of the extreme over-rotating BMPV BH Eq. (4.12). Its $\xi \to 0$ limit is well defined and gives the vacuum with $aDS_3 \times S^2$ geometry. The naive limit $\xi \to \infty$ is singular, though, but can be made regular by using the following observation: under the field rescalings

$$e^a{}_\mu \to \varepsilon^{-1} e^a{}_\mu \,, \qquad V \to \varepsilon^{-1} V \,, \tag{7.1}$$

the action of $N = 1, d = 5$ SUGRA Eq. (2.4) scales homogeneously $S \to \varepsilon^{-3} S$. Then, we can combine this rescaling that transforms solutions into solutions with the reparametrization

$$\chi \to \varepsilon\chi \,, \quad \theta \to \varepsilon\theta \,, \quad t \to \varepsilon t + R_3/2(\cosh\xi\phi - \sinh\xi\varphi) \,, \tag{7.2}$$

and the gauge transformation of the vector field

$$\delta V = \tfrac{\sqrt{3}}{2} R_3(\cosh\xi d\varphi - \sinh\xi d\phi) \,, \tag{7.3}$$

and take the double limit $\xi \to \infty$, $\varepsilon \to 0$, whilst keeping the product $\varepsilon \sinh\xi = 1$. The result is the solution

$$\left\{ \begin{array}{rcl} ds^2 & = & [dt + R_3/4(\chi^2 d\phi - \theta^2 d\varphi)]^2 - (R_3/2)^2[d\chi^2 + \chi^2 d\phi^2 + d\theta^2 + \theta^2 d\varphi^2] \,, \\ V & = & \tfrac{\sqrt{3}}{4} R_3(\theta^2 d\varphi + \chi^2 d\phi) \,, \end{array} \right. \tag{7.4}$$

which is the Gödel solution Eq. (4.16) in polar coordinates with $\lambda = 1/R_3$.

In this limit the number of supersymmetries and symmetries has been preserved. Observe that the gravitino supersymmetry rule Eq. (2.5) $\delta_\epsilon\psi_\mu$ is invariant under the above field rescalings. Actually, the arguments of [49, 50] for the Penrose-Güven limit apply equally to this kind of limit.

In the remaining of this section we are going to apply it to two cases: the Euclidean Robinson-Bertotti solutions of $N = 2_{\pm}, d = 4$ SUGRA Eqs. (5.1),(5.2) and the $aDS_5 \times S^5$ solution of $d = 10$ type IIB SUGRA.

7.1 The limit of the Euclidean Dyonic Robinson-Bertotti solutions

The action and gravitino supersymmetry rule of $N = 2, d = 4$ SUGRA also transform homogeneously under the rescalings Eq. (7.1) and we can use them to take a smooth limit of the Euclidean dyonic Robinson-Bertotti solutions Eqs. (5.1),(5.2), exactly as in the previous case. The calculations are clearly identical and the result is, not surprisingly the flacuum solution Eq. (5.3).

7.2 Boosted Hopf oxidation of $aDS_5 \times S^5$

It is well known that S^5 can be seen as an S^1 fibration over $\mathbb{CP}^2$. In [51], it was observed that the aDS_{2n+1} spacetimes can also be seen as a timelike S^1 fibration over a non-compact version of $\mathbb{CP}^n$, which the authors called $\overline{\mathbb{CP}}^n$. This construction was, then, used to derive solution in the Euclidean IIA theory via timelike T-duality.

The relation of the NH limit of the extreme BMPV BH solutions to the $N = (1, 0), d = 6$ vacuum $aDS_3 \times S^3$, however, shows that in general we before performing dimensional reduction or a T-duality transformation one can mix the S^1 fibers of the sphere and the aDS spacetime by a rotation, if both fibers are spacelike, or by a boost if one is spacelike and the other is timelike.

In the case of $aDS_5 \times S^5$ the only mixing that we can do is a boost on the timelike fiber in the aDS_5 and the spacelike fiber for the S^5. After the boost, we have basically 2 options: T-dualize over the new timelike direction, obtaining a generalization of the solutions discussed in [51], or T-dualize over the new spacelike direction, which will lead to a solution similar to the NH limit of the extreme over-rotating BMPV BH, and that will contain the $aDS_5 \times \mathbb{CP}^2 \times \mathbb{R}$ solution as a special limit.

Let us write the $aDS_5 \times S^5$ solution as

$$\left\{ \begin{array}{rcl} ds^2_{IIB} & = & \big[dt - B(\xi, \overline{\xi})\big)\big]^2 \;-\; \big[dy - A(z, \overline{z})\big]^2 \;-\; d\overline{\mathbb{CP}}^2_{[\xi]} \;-\; d\mathbb{CP}^2_{[z]}\ , \\ G^{(5)} & = & 4\ dt \wedge \omega_{\overline{\mathbb{CP}}^2} \;+\; 4\ dy \wedge \omega_{\mathbb{CP}^2}\ , \end{array} \right. \tag{7.5}$$

where we have introduced $(i, j = 1, 2)$ the metrics

$$
\begin{aligned}
d\mathbb{CP}^2_{[z]} &= \frac{dz^i d\bar{z}^i}{1+|z|^2} - \frac{\bar{z}^i dz^i \; z^j d\bar{z}^j}{(1+|z|^2)^2}\,, & A &= \tfrac{i}{2}\frac{\bar{z}^i dz^i - z^i d\bar{z}^i}{1+|z|^2}\,, \\
d\overline{\mathbb{CP}}^2_{[\xi]} &= \frac{d\xi^i d\bar{\xi}^i}{1-|\xi|^2} + \frac{\xi^i d\bar{\xi}^i \; \bar{\xi}^j d\xi^j}{\left(1-|\xi|^2\right)^2}\,, & B &= \tfrac{i}{2}\frac{\xi^i d\bar{\xi}^i \; - \; \bar{\xi}^i d\xi^i}{1-|\xi|^2}\,,
\end{aligned}
\tag{7.6}
$$

$\omega_{\mathbb{CP}^2}$ and $\omega_{\mathbb{CP}^2}$ being their corresponding volume 4-forms.

Boosting on the (t, y)-plane, with boost parameter β, T-dualizing over y and lifting the solution up to M-theory, we find the solution

$$
\left\{
\begin{aligned}
ds^2_{11} &= \left(dt - \cosh\beta B - \sinh\beta A\right)^2 \; - \; d\overline{\mathbb{CP}}^2_{[\xi]} \; - \; d\mathbb{CP}^2_{[z]} \; - \; d\mathbb{R}^2_{[x]}\,, \\
G &= 4\cosh\beta \; \omega_{\mathbb{CP}^2} - 4\sinh\beta \; \omega_{\overline{\mathbb{CP}}^2} - [\sinh\beta dB + \cosh\beta dA] \wedge dx^1 \wedge dx^2\,.
\end{aligned}
\right.
\tag{7.7}
$$

Now we can take the $\beta \to \infty$ limit using the fact that the action for the bosonic fields of M-theory scales homogeneously under the field rescalings

$$
e^a{}_\mu \to \varepsilon^{-1} e^a{}_\mu\,, \qquad C_{\mu\nu\rho} \to \varepsilon^{-3} C_{\mu\nu\rho}\,. \tag{7.8}
$$

We perform the above rescaling and a change of coordinates

$$
t \to \varepsilon t\,, \quad \xi^i \to \varepsilon \xi^i\,, \quad z^i \to \varepsilon z^i\,, \quad x^i \to \varepsilon x^i\,, \tag{7.9}
$$

and now take the double limit $\beta \to \infty$ $\varepsilon \to 0$ with $\sinh\beta\varepsilon = 1$. It is readily seen that the limiting solution is the $n = 4$ Gödel spacetime that preserves 20 supersymmetries [10].

8 Conclusions

In this paper we have explored the relations between maximally supersymmetric vacua of $d = 4, 5, 6$ Euclidean and Lorentzian SUGRA theories with 8 supercharges and we have found that the maximally supersymmetric graviphoton background of [24, 8] which we have called flacuum solution is related to the maximally supersymmetric Gödel solution by timelike dimensional reduction, a situation that can be generalized to higher-dimensional Gödel solutions of M-theory, for instance, and higher-dimensional flacuum solutions, of Euclidean type II theory, for instance.

We have studied other instances in which Euclidean solutions with vanishing energy-momentum tensor appear, as in the ("double") lightlike dimensional reduction.

Apart from the relations between solutions via spacelike and timelike dimensional reduction and oxidation we have found a limiting procedure similar in may respects to the well-known Penrose-Güven limiting procedure that, instead of giving plane-wave spacetimes gives, at least in the examples considered, Gödel solutions.

The main interest in flacuum solutions, for the moment, seems to be due to the fact that their superalgebras are small deformations of the superalgebra that "defines" the

SUGRA theory in question. All the maximally supersymmetric vacua if a given SUGRA have, in fact, a symmetry superalgebra that can be seen as a deformation of the SUGRA superalgebra in which some central charges are activated and these, together with some momenta, are given non-trivial commutation relations between them and with the supercharges. These new non-vanishing commutators and the central charges vanish for some value of the deformation parameter.

For instance, the (Lorentzian) $N=2, d=4$ Poincaré superalgebra is given by the only on-vanishing anticommutator (in a Majorana basis)

$$\{\mathcal{Q}_{(\alpha i)}, \mathcal{Q}_{(\beta j)}\} = -i\delta_{ij}(\mathcal{C}\gamma^a)_{\alpha\beta}P_a + i\mathcal{C}_{\alpha\beta}\epsilon_{ij}Q + (\mathcal{C}\gamma_5)_{\alpha\beta}\epsilon_{ij}P\,, \tag{8.1}$$

where Q and P are the central charges. All the bosonic charges commute. Now, let us consider different maximally supersymmetric vacua of this theory. Minkowski spacetime has the same superalgebra, with vanishing central charges. The superalgebra of the Robinson-Bertotti solution ($aDS_2 \times S^2$) has the same anticommutator but, now, the six bosonic charges generate an $SO(2,1) \times SO(3)$ Lie algebra

$$\begin{array}{rclrclrcl} [Q,P_1] &=& \frac{1}{R_2}P_3\,, & [Q,P_0] &=& \frac{1}{R_2}P_1\,, & [P_1,P_0] &=& -\frac{1}{R_2}Q\,, \\ [P,P_2] &=& \frac{1}{R_2}P_3\,, & [P,P_3] &=& -\frac{1}{R_2}P_2\,, & [P_2,P_3] &=& \frac{1}{R_2}P\,, \end{array} \tag{8.2}$$

that become again Abelian when the aDS_2 and S^2 radius R_2 goes to infinity. The commutators of the supercharges and these bosonic generators also become nontrivial and have a similar behavior.

Another examples is provided by the KG4 H*pp*-wave solution [6]. The supercharges of its symmetry superalgebra also have the same anticommutator as above, but now the six bosonic charges generate the Heisenberg algebra with non-vanishing commutators

$$\begin{array}{rclrclrcl} [Q,P_1] &=& -\frac{\lambda}{4}P_v\,, & [P,P_2] &=& -\frac{\lambda}{4}P_v\,, & [P_u,P_1] &=& \lambda Q\,, \\ [P_u,P_2] &=& \lambda P\,, & [P_u,Q] &=& -\frac{\lambda}{4}P_1\,, & [P_u,P] &=& -\frac{\lambda}{4}P_2\,, \end{array} \tag{8.3}$$

that also becomes Abelian in the $\lambda \to 0$ limit.

The maximally supersymmetric $aDS_n \times S^m$ vacua of 11- and 10-dimensional supergravities can also be seen as deformations of the Poincaré superalgebra and the same is valid for H*pp*-wave backgrounds.

It is natural to expect that quantum theories defined in these backgrounds satisfy some non-commutative generalization of the Heisenberg algebra.

Acknowledgements

We thank E. Lozano-Tellechea for interesting conversations. T.O. would like to thank M.M. Fernández for her support. The work of T.O. has been supported in part by the Spanish grant BFM2003-01090.

A The detailed dimensional reductions

The dimensional reduction of the Einstein-Hilbert term is identical in the two cases $N = (1,0), \hat{d} = 6$ and $N = 1, \hat{d} = 5$ SUGRA. Further, in both the timelike and spacelike cases we can make the *Ansatz*

$$\left(\hat{e}_{\hat{\mu}}{}^{\hat{a}}\right) = \begin{pmatrix} e_{\mu}{}^{a} & kA_{\mu} \\ \\ 0 & k \end{pmatrix} , \qquad \left(\hat{e}_{\hat{a}}{}^{\hat{\mu}}\right) = \begin{pmatrix} e_{a}{}^{\mu} & -A_{a} \\ \\ 0 & k^{-1} \end{pmatrix} , \tag{A.1}$$

where $A_a = e_a{}^{\mu} A_{\mu}$. All d-dimensional fields with Lorentz indices will be assumed to have been contracted with the d-dimensional Vielbeins.

This *Ansatz* gives the following non-vanishing components of the Ricci rotation coefficients

$$\hat{\Omega}_{abc} = \Omega_{abc} \, , \qquad \hat{\Omega}_{ab\sharp} = \tfrac{1}{2}\alpha_{\hat{d}} F_{ab} \, , \qquad \hat{\Omega}_{a\sharp\sharp} = \tfrac{1}{2}\alpha_{\hat{d}} \partial_a \log k \, , \tag{A.2}$$

and of the the spin connection $\hat{\omega}_{\hat{a}\hat{b}\hat{c}}$

$$\begin{array}{rclcrcl} \hat{\omega}_{abc} & = & \omega_{abc} \, , & \qquad & \hat{\omega}_{ab\sharp} & = & -\tfrac{1}{2}\alpha_{\hat{d}} k F_{ab} \, , \\ \\ \hat{\omega}_{\sharp bc} & = & \tfrac{1}{2}\alpha_{\hat{d}} k F_{bc} \, , & & \hat{\omega}_{\sharp b \sharp} & = & \alpha \partial_b \log k \, , \end{array} \tag{A.3}$$

where

$$F_{ab} = e_a{}^{\mu} e_b{}^{\nu} F_{\mu\nu} \, , \qquad F_{\mu\nu} = 2\partial_{[\mu} A_{\nu]} \, , \tag{A.4}$$

(which we will also denote by $F(A)$ in presence of other vector fields) is the KK vector field strength. The standard procedure gives the KK-frame action

$$\hat{S} = \int d^{\hat{d}}\hat{x} \, \sqrt{|\hat{g}|} \, \hat{R} = \int d^{\hat{d}-1}x \sqrt{|g|} \, k \left[R + \tfrac{1}{4}\alpha_{\hat{d}} k^2 F^2 \right] \, . \tag{A.5}$$

The sign of the Maxwell term in the timelike case is unconventional because we would get the opposite sign by Wick-rotating the standard Lorentzian Maxwell action. However, it cannot be deemed as "wrong" since there is no propagation, no concepts of motion or energy in Euclidean space. On the other hand, if we Wick-rotated a pseudovector field we would get precisely the above sign.

A.1 $N = (1,0), \hat{d} = 6$ SUGRA

In the reduction of the action of $N = (1,0), \hat{d} = 6$ SUGRA Eq. (2.1) to $d = 5$ we have to take into account the anti-selfduality constraint, which can be imposed on the action *after* dimensional reduction, just as in the reduction of $N = 2B, d = 10$ supergravity on a circle [44]. Thus, we first perform the naive dimensional reduction of Eq. (2.1).

We can use our previous results for the reduction of the Einstein-Hilbert term and we only need to reduce the 2-form kinetic term. The reduction of $\hat{H}^{-}$ gives a 3- and a 2-form field strengths

$$
\begin{aligned}
H_{abc} &= \hat{H}^{-}_{abc}\,, & H &= 3\left[\partial B - \tfrac{1}{2}AF(B) - \tfrac{1}{2}BF(A)\right]\,, \\
F_{ab}(B) &= k^{-1}\hat{H}^{-}_{ab\sharp}\,, & F(B) &= 2\partial B\,,
\end{aligned}
\tag{A.6}
$$

where

$$
\hat{B}_{\mu\nu} = B_{\mu\nu} - A_{[\mu}B_{\nu]}\,, \qquad B_{\mu} = \hat{B}_{\mu\underline{\sharp}}\,. \tag{A.7}
$$

The anti-selfduality constraint becomes

$$
{}^{\star}H = -k^{-1}F(B) \;\;\Leftrightarrow\;\; H = \alpha_6 k^{-1\star}F(B)\,. \tag{A.8}
$$

We immediately get

$$
S = \int d^5x\sqrt{|g|}\,k\,[R + \tfrac{1}{4}\alpha_6 k^2F^2(A) + \tfrac{1}{8}\alpha_6 k^{-2}F^2(B) + \tfrac{1}{24}H^2]\,. \tag{A.9}
$$

As in Ref. [44], we Poincaré-dualize the 2-form into a third vector field C_μ and then we identify it with B_μ in the action, that is left with the metric and the unconstrained vector fields A_μ and B_μ and takes the form

$$
S = \int d^5x\sqrt{|g|}\,k\,[R + \tfrac{1}{4}\alpha_6 k^2F^2(A) + \tfrac{1}{4}\alpha_6 k^{-2}F^2(B) - \alpha_6\tfrac{\epsilon}{8\sqrt{|g|}}k^{-1}F(A)F(B)B]\,. \tag{A.10}
$$

The truncation to pure supergravity involves setting to zero the scalar $k = 1$ consistently, i.e. in such a way that its equation of motion is always satisfied. The k equation of motion with $k = 1$ (upon use of Einstein's equation) implies the constraint

$$
F^2(B) = 2F^2(A)\,. \tag{A.11}
$$

We also have to set to zero the gaugino λ and the matter vector field W_μ, which must be a combination of A_μ and B_μ related to λ by supersymmetry. Thus, to identify W_μ, we have to analyze the reduction and truncation of the gravitino supersymmetry transformation rule Eq. (2.2).

The 6- and 5-dimensional gamma matrices are related by

$$
\begin{aligned}
\hat{\gamma}^{a} &= \gamma^{a}\otimes\sigma^{1}\,, \\
\hat{\gamma}^{\sharp} &= \mathbb{I}\otimes\alpha_6^{1/2}\sigma^{2} \equiv \hat{\gamma}^{\sharp}\,, \\
\hat{\gamma}_{7} &= \hat{\gamma}_0\cdots\hat{\gamma}_5 = \mathbb{I}\otimes\sigma^{3}\,,
\end{aligned}
\tag{A.12}
$$

where the γ^as Lorentzian or Euclidean 5-dimensional gamma matrices satisfying $\gamma_0\cdots\gamma_4 = \mathbb{I}$ and $\gamma_1\cdots\gamma_5 = i\mathbb{I}$.

Using this relation, the decompositions of the spin connection and 3-form field strength, the anti-selfduality constraint Eq. (A.8), the chirality of $\hat{\epsilon}^{+}$ and assuming that the supersymmetry parameter $\hat{\epsilon}^{+}$ is independent of $x^{\sharp}$ and setting $k = 1$, we find

$$\begin{aligned} \delta_{\hat{\epsilon}^{+}} \hat{\psi}_{a}^{+} &= \left\{ \nabla_{a} - \tfrac{i}{4} \alpha_{6}^{3/2} \not{F}_{a}(A) + \tfrac{i}{8\sqrt{2}} \alpha_{6}^{1/2} \not{F}(B) \gamma_{a} \right\} \epsilon \,, \\ \delta_{\hat{\epsilon}^{+}} \hat{\psi}_{\sharp}^{+} &= \tfrac{-1}{8\sqrt{2}} \left\{ \sqrt{2} \alpha_{6} \not{F}(A) + \not{F}(B) \right\} \epsilon \,, \end{aligned} \tag{A.13}$$

where $\frac{1}{2}(1+\sigma^{3})\hat{\epsilon}^{+} = \epsilon$.

$\hat{\psi}_{a}^{+}$ is to be identified with the 5-dimensional gravitino $\frac{1}{2}(1+\sigma^{3})\hat{\psi}_{a}^{+} = \psi_{a}$. $\hat{\psi}_{\sharp}^{+}$ only transforms into vectors and it is to be identified with the gaugino $\frac{1}{2}(1+\sigma^{3})\hat{\psi}_{\sharp}^{+} \equiv \lambda$. This, in turn, implies that the combination of vector field strengths in the r.h.s. is proportional to $F(W)$ (for $k = 1$). The field strengths of V and W must be an $SO(2)$ rotation of those of A and B to preserve the canonical normalization and diagonal form of the kinetic terms in the action. The right combination is that in which λ only transforms into W and not into V:

$$\begin{aligned} F(V) &\equiv \tfrac{1}{\sqrt{3}} F(A) - \sqrt{\tfrac{3}{2}} \alpha_{6} F(B) \,, \\ F(W) &\equiv \sqrt{\tfrac{2}{3}} \alpha_{6} F(A) + \tfrac{1}{\sqrt{3}} F(B) \,. \end{aligned} \tag{A.14}$$

Setting $W_{\mu} = 0$ (to make consistent $\lambda = 0$) implies the constraint

$$\sqrt{2} \alpha_{6} F(A) = -F(B) \,, \tag{A.15}$$

which implies the constraint Eq. (A.11).

Therefore, we can truncate the vector multiplet setting consistently $k = 1, \lambda = 0, W_{\mu} = 0$. The result is the action of pure Lorentzian or Euclidean $N = 1, d = 5$ SUGRA Eq. (2.4) and its gravitino supersymmetry transformation law Eq. (2.5).

A.2 $N = 1, \hat{d} = 5$ SUGRA

The reduction of the Einstein-Hilbert term in the action Eq. (2.4) goes as in the previous case. The reduction of the Maxwell field $\hat{V}_{\hat{\mu}}$ gives rise to a d-dimensional vector field that we denote by B_{μ} and a scalar l:

$$\hat{V}_{\underline{\sharp}} = l \,, \qquad \hat{V}_{\mu} = B_{\mu} + l A_{\mu} \,. \tag{A.16}$$

As for the field strength, we define $F_{\mu\nu}(B)$ and the combination $\mathcal{G}_{ab} = \hat{F}_{ab}$

$$F_{\mu\nu}(B) = 2\partial_{[\mu} B_{\nu]} \,, \qquad \mathcal{G} = F(B) + l F(A) \,, \tag{A.17}$$

where $F(A)$ is the KK vector field strength. Taking into account

$$\hat{F}_{a\sharp} = k^{-1}\partial_a l \tag{A.18}$$

we find that the reduction of the Einstein-Hilbert and Maxwell terms of the action Eq. (2.4) gives

$$\int d^4x\sqrt{|g|}\; k\left[R + \tfrac{1}{2}\alpha_5\alpha_6 k^{-2}(\partial l)^2 + \tfrac{1}{4}\alpha_5 k^2 F^2(A) + \tfrac{1}{4}\alpha_6 \mathcal{G}^2\right] . \tag{A.19}$$

The reduction of the Chern-Simons term is identical for both the timelike and spacelike cases, using $\hat{\epsilon}^{abcd\sharp} = \epsilon^{abcd}$, and gives after integration by parts the $d = 4$ action

$$\begin{aligned} S \;=\; & \int d^4x\sqrt{|g|}\, k\Big\{R + \tfrac{1}{2}\alpha_5\alpha_6 k^{-2}(\partial l)^2 + \tfrac{1}{4}\alpha_5 k^2 F^2(A) + \tfrac{1}{4}\alpha_6\mathcal{G}^2 \\ & \qquad -\tfrac{1}{4\sqrt{3}}\alpha_6 k^{-1} l \tfrac{\epsilon}{\sqrt{|g|}}[\mathcal{G} + 2A\partial l]^2\Big\} . \end{aligned} \tag{A.20}$$

This theory is pure (Euclidean $\alpha_6\alpha_5 = -1$ or Lorentzian $\alpha_6\alpha_5 = +1$) $N = 2, d = 4$ SUGRA theory coupled to a "matter" vector supermultiplet that contains a vector and the two scalars k and l [30]. The matter and supergravity vector fields are combinations of the two vectors A, V. To identify the mater vector field we can use the fact that eliminating a matter supermultiplet is always a consistent truncation. The equations of motion of k and l after setting $k = 1$ and $l = 0$ give the constraints

$$\left\{\begin{aligned} 3\alpha_5 F^2(A) + \alpha_6 F^2(B) &= 0\,, \\ \left[\sqrt{3}F(A) - {}^\star F(B)\right] F(B) &= 0\,. \end{aligned}\right. \tag{A.21}$$

The second constraint is solved by

$$\sqrt{3}F(A) - {}^\star F(B) = 0, \tag{A.22}$$

which also solves the first[20]. This implies that the matter vector field that we denote by W_μ is proportional to the combination $\sqrt{3}F(B) - {}^\star F(B)$ or to its Hodge dual, at least for the $k = 1,\ l = 0$ case. The orthogonal combination will be the field strength of the supergravity vector that we will denote by V_μ.

To fully identify the supergravity and matter vector field strengths we have to check the consistency of the truncation from the supersymmetric point of view, that is, in the gravitino supersymmetry transformation rule Eq. (2.5) (adding hats everywhere). Assuming that the supersymmetry parameter $\hat{\epsilon}$ is independent of the internal coordinate so it can be identified with the 4-dimensional one ϵ we get and setting from now on $k = 1$ and $l = 0$

---

[20] Observe that we have $({}^\star F)^2 = -\alpha_6\alpha_5 F^2$ and ${}^{\star\star}F = -\alpha_6\alpha_5 F$ because $\alpha_5\alpha_6 = +1$ corresponds to Lorentzian signature and $\alpha_5\alpha_6 = -1$ to Euclidean signature.

$$\begin{aligned} \delta_\epsilon \hat{\psi}_a &= \Big\{\nabla_a - \tfrac{1}{4}\alpha_5 \not{F}_a(A)\gamma^\sharp - \tfrac{i}{8\sqrt{3}}\alpha_6^{3/2}[\not{F}(B)\gamma_a + 2 \not{F}_a(B)]\Big\}\,\epsilon\,, \\ \delta_\epsilon \hat{\psi}_\sharp &= -\tfrac{1}{8\sqrt{3}}\alpha_5 \left\{\sqrt{3} \not{F}(A) + \not{F}(B)\gamma^\sharp\right\}\epsilon\,, \end{aligned} \tag{A.23}$$

where $\gamma^a = \hat{\gamma}^a$ with $a = 0,1,2,3$ in the Lorentzian case $\alpha_5\alpha_6 = +1$, and $a = 1,2,3,4$ in the Euclidean case $\alpha_5\alpha_6 = -1$, and $\gamma^\sharp = \hat{\gamma}^\sharp$ is a matrix proportional to the 4-dimensional γ_5. It can be shown that

$$\begin{aligned} \not{F}_a\gamma^\sharp &= -i\alpha_6^{1/2}\left(\tfrac{1}{2}{}^\star\!\not{F}\gamma_a - {}^\star\!\not{F}_a\right)\,, \\ \not{F}\gamma^\sharp &= i\alpha_6^{1/2}\,{}^\star\!\not{F}\,, \end{aligned} \tag{A.24}$$

and using these identities we can rewrite the gravitino supersymmetry transformation rule as follows:

$$\begin{aligned} \delta_\epsilon \hat{\psi}_a &= \Big\{\nabla_a - \tfrac{i}{8}\alpha_6^{3/2}\Big[-\alpha_5\alpha_6{}^\star\!\not{F}(A) + \tfrac{1}{\sqrt{3}} \not{F}(B)\Big]\gamma_a + \tfrac{1}{4}\alpha_6^{3/2}\Big[-\alpha_5\alpha_6{}^\star\!\not{F}_a(A) - \tfrac{1}{\sqrt{3}} \not{F}_a(B)\Big]\Big\}\,\epsilon\,, \\ \delta_\epsilon \hat{\psi}_\sharp &= -\tfrac{1}{8}\alpha_5\Big[\not{F}(A) - \tfrac{1}{\sqrt{3}}{}^\star\!\not{F}(B)\Big]\,\epsilon\,, \end{aligned} \tag{A.25}$$

The components $\hat{\psi}_a$ clearly become the 4-dimensional gravitino ψ_a. The internal component of the gravitino $\hat{\psi}_\sharp$ only transforms into a combination vector field strengths and, thus, it can be identified with the gaugino λ of the matter vector multiplet. The combination of vector field strengths is proportional to the matter vector field strength $F(W)$ (for $k = 1$ and $l = 0$ only), in agreement with the constraint Eq. (A.22) found in the truncation of the action. The normalization factor and the definition of the supergravity vector field strength $F(V)$ are found by imposing the diagonalization and correct normalization of the energy-momentum tensors of the vectors in the Einstein equation: the r.h.s. of the Einstein equation contains

$$-\tfrac{1}{2}\alpha_5 T_{\mu\nu}(A) - \tfrac{1}{2}\alpha_6 T_{\mu\nu}(B)\,, \tag{A.26}$$

where $T_{\mu\nu}$, the energy-momentum tensor of a vector field in four Euclidean or Lorentzian dimensions, is usually written in the form

$$T_{\mu\nu} = F_\mu{}^\rho F_{\nu\rho} - \tfrac{1}{2}g_{\mu\nu}F^2\,, \tag{A.27}$$

but can be rewritten in the more useful form

$$T_{\mu\nu} = \tfrac{1}{2}\left[F_\mu{}^\rho F_{\nu\rho} + \alpha_5\alpha_6{}^\star F_\mu{}^{\rho\star} F_{\nu\rho}\right]\,. \tag{A.28}$$

This still leaves some ambiguities in the identification of the supergravity vector field, but they are irrelevant when $W = 0$ and lead to a unique relation

$$F(V) = 2\alpha_5\alpha_6{}^\star F(A)\,. \tag{A.29}$$

Now we can eliminate consistently the gaugino λ, the matter vector field W_μ and scalars k and l, getting the bosonic action[21] and gravitino supersymmetry transformation rule which are given in Eq. (3.1).

References

[1] J. P. Gauntlett, J. B. Gutowski, C. M. Hull, S. Pakis and H. S. Reall, Class. Quant. Grav. **20** (2003) 4587 [arXiv:hep-th/0209114].

[2] J. B. Gutowski, D. Martelli and H. S. Reall, Class. Quant. Grav. **20** (2003) 5049 [arXiv:hep-th/0306235].

[3] A. Chamseddine, J. Figueroa-O'Farrill and W. Sabra, arXiv:hep-th/0306278.

[4] M. M. Caldarelli and D. Klemm, JHEP **0309** (2003) 019 [arXiv:hep-th/0307022].

[5] E. Lozano-Tellechea, P. Meessen and T. Ortín, Class. Quant. Grav. **19** (2002) 5921 [arXiv:hep-th/0206200].

[6] J. Kowalski-Glikman, Phys. Lett. B **150** (1985) 125.

[7] I. Robinson, *Bull. Acad. Polon. Sci.* **7**(1959), 351; B. Bertotti, *Phys. Rev.* **116** (1959) 1331.

[8] N. Berkovits and N. Seiberg, JHEP **0307** (2003) 010 [arXiv:hep-th/0306226].

[9] B. Fiol, C. Hofman and E. Lozano-Tellechea, arXiv:hep-th/0312209.

[10] T. Harmark and T. Takayanagi, Nucl. Phys. B **662** (2003) 3 [arXiv:hep-th/0301206].

[11] M. M. Caldarelli and D. Klemm, arXiv:hep-th/0310081.

[12] K. Gödel, Rev. Mod. Phys. **21** (1949) 447.

[13] G. T. Horowitz and A. A. Tseytlin, Phys. Rev. D **51** (1995) 2896 [arXiv:hep-th/9409021].

[14] J. G. Russo and A. A. Tseytlin, Nucl. Phys. B **448** (1995) 293 [arXiv:hep-th/9411099].

[15] M. Gurses, A. Karasu and O. Sarioglu, arXiv:hep-th/0312290

[16] E. G. Gimon and A. Hashimoto, Phys. Rev. Lett. **91** (2003) 021601 [arXiv:hep-th/0304181].

---

[21] The bosonic action cannot be found by performing the above field redefinitions and has to bes constructed so as to reproduce the truncated equations of motion.

[17] D. Brecher, U. H. Danielsson, J. P. Gregory and M. E. Olsson, JHEP **0311** (2003) 033 [arXiv:hep-th/0309058].

[18] M. Blau, P. Meessen and M. O'Loughlin, JHEP **0309** (2003) 072 [arXiv:hep-th/0306161].

[19] D. Brace, C. A. R. Herdeiro and S. Hirano, arXiv:hep-th/0307265.

[20] E. K. Boyda, S. Ganguli, P. Horava and U. Varadarajan, Phys. Rev. D **67** (2003) 106003 [arXiv:hep-th/0212087].

[21] N. Drukker, B. Fiol and J. Simon, Phys. Rev. Lett. **91** (2003) 231601 [arXiv:hep-th/0306057].

[22] Y. Hikida and S. J. Rey, Nucl. Phys. B **669** (2003) 57 [arXiv:hep-th/0306148].

[23] D. Brecher, P. A. DeBoer, D. C. Page and M. Rozali, JHEP **0310** (2003) 031 [arXiv:hep-th/0306190].

[24] N. Seiberg, JHEP **0306** (2003) 010 [arXiv:hep-th/0305248].

[25] H. Ooguri and C. Vafa, Adv. Theor. Math. Phys. **7** (2003) 53 [arXiv:hep-th/0302109].

[26] S. B. Giddings and A. Strominger, Phys. Lett. B **230** (1989) 46.

[27] G. W. Gibbons, M. B. Green and M. J. Perry, Phys. Lett. B **370** (1996) 37 [arXiv:hep-th/9511080].

[28] H. Nishino and E. Sezgin, Nucl. Phys. B **278** (1986) 353.

[29] E. Cremmer, LPTENS 80/17 *Invited paper at the Nuffield Gravity Workshop, Cambridge, Eng., Jun 22 - Jul 12, 1980*

[30] A. H. Chamseddine and H. Nicolai, Phys. Lett. B **96** (1980) 89.

[31] P. Meessen, Phys. Rev. D **65** (2002) 087501 [arXiv:hep-th/0111031].

[32] G. W. Gibbons, G. T. Horowitz and P. K. Townsend, Class. Quant. Grav. **12** (1995) 297 [arXiv:hep-th/9410073].

[33] J. C. Breckenridge, R. C. Myers, A. W. Peet and C. Vafa, Phys. Lett. B **391** (1997) 93 [arXiv:hep-th/9602065].

[34] R. Kallosh, A. Rajaraman and W. K. Wong, Phys. Rev. D **55** (1997) 3246 [arXiv:hep-th/9611094].

[35] J. P. Gauntlett, R. C. Myers and P. K. Townsend, Class. Quant. Grav. **16** (1999) 1 [arXiv:hep-th/9810204].

[36] K. P. Tod, Phys. Lett. B **121** (1983) 241.

[37] A. A. Belavin, A. M. Polyakov, A. S. Shvarts and Y. S. Tyupkin, Phys. Lett. B **59** (1975) 85.

[38] J. M. Figueroa-O'Farrill, Class. Quant. Grav. **16** (1999) 2043 [arXiv:hep-th/9902066].

[39] T. Ortín, Class. Quant. Grav. **19** (2002) L143 [arXiv:hep-th/0206159].

[40] N. Alonso-Alberca, E. Lozano-Tellechea and T. Ortín, arXiv:hep-th/0208158.

[41] A. González-Arroyo, arXiv:hep-th/9807108.

[42] C. Hull, arXiv:hep-th/0305039.

[43] G. Papadopoulos and D. Tsimpis, Class. Quant. Grav. **20** (2003) L253 [arXiv:hep-th/0307127].

[44] P. Meessen and T. Ortín, Nucl. Phys. B **541** (1999) 195 [arXiv:hep-th/9806120].

[45] A. Trautman, Int. J. Theor. Phys. **16** (1977) 561.

[46] B. Julia and H. Nicolai, Nucl. Phys. B **439** (1995) 291 [arXiv:hep-th/9412002].

[47] R. Penrose, in *Differential Geometry and Relativity*, Reidel, Dordrecht (1976) 271-275.

[48] R. Gueven, Phys. Lett. B **482** (2000) 255 [arXiv:hep-th/0005061].

[49] M. Blau, J. Figueroa-O'Farrill, C. Hull and G. Papadopoulos, arXiv:hep-th/0201081.

[50] M. Blau, J. Figueroa-O'Farrill and G. Papadopoulos, arXiv:hep-th/0202111.

[51] C. N. Pope, A. Sadrzadeh and S. R. Scuro, "*Timelike Hopf duality and type IIA* string solutions*", Class. Quant. Grav. **17** (2000) 623 [arXiv:hep-th/9905161].